

---

# Some properties of realizability models

Jean-Louis Krivine

PPS Group, University Paris-Diderot, CNRS

[krivine@pps.univ-paris-diderot.fr](mailto:krivine@pps.univ-paris-diderot.fr)

Chambéry, June 2012

---

## Outline

We give some general properties of classical realizability and we look at some particular models :

- True *arithmetical formulas*, and even true  $\Pi_1^1$  *formulas* are realized ; thus, realizability models cannot give undecidability results in arithmetic.
- A model is given by forcing iff its Boolean algebra  $\mathbb{B}$  is *trivial*.
- We build models in which  $\mathbb{B}$  is *non trivial* and *finite*.
- Following T. Ehrhard and T. Streicher, the usual models of lambda-calculus have, in fact, a structure of realizability algebra.

Therefore, they give rise to realizability models of ZF.

We study a simple case, in which  $\mathbb{B}$  is *non trivial* and *integers are preserved*.

---

## A game on first order formulas

We consider first order formulas written with :

$\rightarrow, \forall, \top, \perp, \neq$ , predicate constants, function symbols for recursive functions.

A 1st order formula has the form  $\forall \vec{x}[\Phi_1, \dots, \Phi_n \rightarrow A]$  where  $\Phi_1, \dots, \Phi_n$  are 1st order formulas and  $A$  is atomic (i.e.  $Rt_1 \dots t_k$  or  $t_0 \neq t_1$  or  $\top$  or  $\perp$ ).

In the following, we only consider *closed* 1st order formulas.

The atomic closed formula  $t_0 \neq t_1$  is interpreted as  $\top$  (resp.  $\perp$ ) if it is true (resp. false) in  $\mathbb{N}$ .

We define a game with two players :  $\exists$  (the *client*) and  $\forall$  (the *server*).

At each step, the *position* is a sequent  $\mathcal{U} \vdash \mathcal{A}$  with closed 1st order formulas ; the formulas of  $\mathcal{A}$  are atomic and  $\perp \in \mathcal{A}$  ;  $\mathcal{U}$  and  $\mathcal{A}$  increase at each step.

The game starts with a sequent  $\mathcal{U}_0 \vdash \mathcal{A}_0$ .

---

A move in this game is as follows :

Player  $\exists$  chooses  $\Psi \in \mathcal{U}$ ,  $\Psi = \forall \vec{y}[\Phi_1(\vec{y}), \dots, \Phi_n(\vec{y}) \rightarrow B(\vec{y})]$

and  $\vec{j} \in \mathbb{N}^l$  such that  $B(\vec{j}) \in \mathcal{A}$  (if this is impossible, then  $\exists$  has lost).

Player  $\forall$  chooses a formula  $\Phi \in \mathcal{V} = \{\Phi_1(\vec{j}), \dots, \Phi_n(\vec{j})\}$ ,

$\Phi \equiv \forall \vec{x}[\Psi_1(\vec{x}), \dots, \Psi_m(\vec{x}) \rightarrow A(\vec{x})]$  ;  $\forall$  chooses also  $\vec{i} \in \mathbb{N}^k$ .

The atomic formula  $A(\vec{i})$  must not be  $\top$  (otherwise,  $\forall$  has lost).

Then  $\Psi_1(\vec{i}), \dots, \Psi_m(\vec{i})$  *are added* to  $\mathcal{U}$  and  $A(\vec{i})$  *is added* to  $\mathcal{A}$ .

$\exists$  wins iff  $\forall$  cannot play at some step

(every formula of  $\mathcal{V}$  ends with  $\top$ , in particular if  $\mathcal{V} = \emptyset$ ).

In fact, player  $\forall$  tries to build a model over  $\mathbb{N}$  in which

the formula  $\mathcal{V}_0 = \bigwedge \mathcal{U}_0 \rightarrow \bigvee \mathcal{A}_0$  is false, and  $\exists$  tries to avoid this :

---

**Theorem.** i) Any model  $\mathcal{M}$  over  $\mathbb{N}$  s.t.  $\mathcal{M} \not\models \mathcal{V}_0$  gives a winning strategy for  $\forall$ .

ii) There exists a “trivial” strategy for the player  $\exists$  such that each play  $\exists$  loses using it, gives a model  $\mathcal{M}$  over  $\mathbb{N}$ ,  $\mathcal{M} \not\models \mathcal{V}_0$ .

i) We define a strategy for  $\forall$  such that, at each step :  
every formula of  $\mathcal{U}$  (resp.  $\mathcal{A}$ ) is true (resp. false) in  $\mathcal{M}$ .

This is true at the beginning of the game.

Then  $\exists$  chooses  $\Psi \in \mathcal{U}$ ,  $\Psi = \forall \vec{y}[\Phi_1(\vec{y}), \dots, \Phi_n(\vec{y}) \rightarrow B(\vec{y})]$  and  $\vec{j} \in \mathbb{N}^l$   
such that  $B(\vec{j}) \in \mathcal{A}$ . Therefore,  $\mathcal{M} \models \neg B(\vec{j})$  and  $\mathcal{M} \models \Psi$ .

Thus,  $\forall$  can choose  $\Phi \in \mathcal{V} = \{\Phi_1(\vec{j}), \dots, \Phi_n(\vec{j})\}$  s.t.  $\mathcal{M} \models \neg \Phi$ .

Let  $\Phi = \forall \vec{x}[\Psi_1(\vec{x}), \dots, \Psi_m(\vec{x}) \rightarrow A(\vec{x})]$ .

Then  $\forall$  can choose  $\vec{i} \in \mathbb{N}^k$  s.t.  $\mathcal{M} \models \Psi_1(\vec{i}), \dots, \Psi_m(\vec{i})$  and  $\neg A(\vec{i})$ .

Finally  $\Psi_1(\vec{i}), \dots, \Psi_m(\vec{i})$  are added to  $\mathcal{U}$  and  $A(\vec{i})$  to  $\mathcal{A}$ .

Thus  $\mathcal{U}$  and the negation of formulas of  $\mathcal{A}$  remain true in  $\mathcal{M}$ .

---

ii) Here is the “trivial” strategy for  $\exists$  :

*fix an enumeration of all ordered pairs  $\langle \Psi, \vec{j} \rangle$  ( $\Psi$  is a closed formula,  $\vec{j} \in \mathbb{N}^l$ ).*

*At each step,  $\exists$  chooses the first allowed pair  $\langle \Psi, \vec{j} \rangle$ , not chosen before.*

Suppose  $\exists$  loses some play with this strategy. Let  $\mathcal{M}$  be the model which satisfies exactly the closed atomic formulas never put in  $\mathcal{A}$  during this play.

A pair  $\langle \Psi, \vec{j} \rangle$  is called *acceptable* if  $\Psi$  is put in  $\mathcal{U}$  and  $B(\vec{j})$  in  $\mathcal{A}$  at some step (not necessarily the same) where  $B(\vec{j})$  is the final atom of  $\Psi$ .

Every acceptable pair is effectively played by  $\exists$  at some step :

namely when every acceptable pair strictly less than it has been played.

We prove, by induction, that  $\mathcal{M}$  satisfies *every formula  $\Psi$  which is put in  $\mathcal{U}$*  and the negation of *every formula  $\Phi$  chosen by  $\forall$*  during the play.

---

**Proof for  $\Psi$ .** The result is clear if  $\Psi$  is atomic because, if  $\Psi$  is both in  $\mathcal{U}$  and  $\mathcal{A}$  then  $\langle \Psi, \emptyset \rangle$  is acceptable and thus will be chosen by  $\exists$ ; then  $\exists$  wins.

Otherwise, let  $\Psi = \forall \vec{y}[\Phi_1(\vec{y}), \dots, \Phi_n(\vec{y}) \rightarrow B(\vec{y})]$ . We must show that  $\mathcal{M} \models \Phi_1(\vec{j}), \dots, \Phi_n(\vec{j}) \rightarrow B(\vec{j})$  for every  $\vec{j} \in \mathbb{N}^k$ .

This is clear if  $B(\vec{j})$  is never put in  $\mathcal{A}$ , because  $\mathcal{M} \models B(\vec{j})$ .

Otherwise,  $\langle \Psi, \vec{j} \rangle$  is acceptable and is chosen by  $\exists$  at some step.

Then  $\mathcal{V} = \{\Phi_1(\vec{j}), \dots, \Phi_n(\vec{j})\}$  and  $\Phi_1(\vec{j})$ , for instance, is chosen by  $\forall$ .

By induction hypothesis, we have  $\mathcal{M} \models \neg \Phi_1(\vec{j})$ , which gives the result.

**Proof for  $\Phi$ .** Let  $\Phi = \forall \vec{x}[\Psi_1(\vec{x}), \dots, \Psi_m(\vec{x}) \rightarrow A(\vec{x})]$ ;  $\forall$  chooses  $\vec{i}$  and puts  $A(\vec{i})$  in  $\mathcal{A}$  and  $\Psi_1(\vec{i}), \dots, \Psi_m(\vec{i})$  in  $\mathcal{U}$ . By induction hypothesis,  $\mathcal{M} \models \Psi_1(\vec{i}), \dots, \Psi_m(\vec{i})$ ; and, by definition,  $\mathcal{M} \not\models A(\vec{i})$ . Thus  $\mathcal{M} \models \neg \Phi$ .

It follows that  $\mathcal{M} \not\models \mathcal{V}_0$  since  $\mathcal{M} \models \mathcal{U}_0$  and  $\mathcal{M} \models \neg A$  for  $A \in \mathcal{A}_0$ .

QED

---

## Well founded recursive relations

Let  $f : \mathbb{N}^2 \rightarrow \{0, 1\}$  be arbitrary. The predicate  $f(x, y) = 1$  is well founded iff the formula  $\forall X \forall z \{ \forall x [ \forall y (f(x, y) = 1 \rightarrow Xy) \rightarrow Xx ] \rightarrow Xz \}$  is true in  $\mathbb{N}$ .

We show that, in this case, this formula is even *realized*.

**Theorem.** If the predicate  $f(x, y) = 1$  is well founded, then

$\Upsilon \Vdash \forall X \forall z \{ \forall x [ \forall y (f(x, y) = 1 \mapsto Xy) \rightarrow Xx ] \rightarrow Xz \}$ .

Let  $t \Vdash \forall x [ \forall y (f(x, y) = 1 \mapsto Xy) \rightarrow Xx ]$  and  $n \in \mathbb{N}$ ; we show by induction on  $n$ , following the well founded predicate " $f(x, y) = 1$ ", that  $\Upsilon t \Vdash Xn$ .

Since  $\Upsilon t \star \pi > t \star \Upsilon t \bullet \pi$ , it suffices to show that  $\Upsilon t \Vdash \forall y (f(n, y) = 1 \mapsto Xy)$

i.e.  $\Upsilon t \Vdash f(n, p) = 1 \mapsto Xp$ . This is trivial if  $f(n, p) \neq 1$

and this follows from the induction hypothesis if  $f(n, p) = 1$ .

Thus, if  $\pi \in \Vdash Xn \Vdash$ , we have  $t \star \Upsilon t \bullet \pi \in \perp$  and therefore  $\Upsilon \star t \bullet \pi \in \perp$ .

QED

This shows that a *recursive* well founded predicate on integers is also well founded *in every realisability model*.

---



---

## True $\Pi_1^1$ formulas

A  $\Pi_1^1$  formula is of the form  $F \equiv \forall \vec{X} \Phi[\vec{X}]$  where  $\Phi$  is a 1st order formula written with the function symbols  $0, 1, +, \times$  and the predicate symbols  $\neq, \vec{X}$ .

**Theorem.** If  $F$  is a true  $\Pi_1^1$  formula, then  $F^{\text{int}}$  is realized.

This shows, in particular, that the integers of any realizability model are *elementary equivalent* to the integers of the ground model.

It is not possible to show the independence of some *arithmetical* (and even  $\Pi_1^1$ ) formula by means of realizability models.

Open problems : What about  $\Sigma_1^1$  (or higher) formulas ?

Are the *constructible universes* of the ground model and the realizability model elementarily equivalent ? This is (trivially) true in the case of forcing.

---

**Proof.** Fix a recursive enumeration of closed formulas and also of sequents  $\mathcal{U} \vdash \mathcal{A}$ .

Let  $F \equiv \forall \vec{X} \neg \Phi[\vec{X}]$  be a *true*  $\Pi_1^1$  formula.

The meaning of  $F$  is that the 1st order formula  $\Phi \rightarrow \perp$  has no model.

Thus, *the “trivial” strategy for  $\exists$  is winning*

in the game which starts with the sequent  $\Phi \vdash \perp$ .

Now, let  $f(x, y) = 1$  be the recursive predicate which says that

*$x, y$  are (numbers of) successive positions chosen by  $\forall$  such that, between them,  $\exists$  has applied (once) the trivial strategy.*

This strategy is winning for  $\exists$  iff each play is finite, i.e. iff *the predicate  $f(x, y) = 1$  is well founded.*

Now, by the above theorem, we obtain :

$Y \Vdash \forall X \{ \forall x [ \forall y ( f(x, y) = 1 \mapsto Xy ) \rightarrow Xx ] \rightarrow \forall x Xx \}$ .

But we have just proved that : *“ $f(x, y) = 1$  is well founded”  $\rightarrow F$ .*

Let  $\theta$  be a proof-like term associated with this proof. Then  $\theta Y \Vdash F$ .

**QED**

---

## The case of arithmetical formulas

An arithmetical formula is of the form

$$\forall x_1 \exists y_1 \dots \forall x_n \exists y_n (f(x_1, y_1, \dots, x_n, y_n) \neq 0)$$

where  $f : \mathbb{N}^{2n} \rightarrow \{0, 1\}$  is recursive.

**Theorem.** Let  $f : \mathbb{N}^{2n} \rightarrow \{0, 1\}$  be an arbitrary function, such that

$\forall x_1 \exists y_1 \dots \forall x_n \exists y_n (f(x_1, y_1, \dots, x_n, y_n) \neq 0)$  is true in  $\mathbb{N}$ . Then

$\forall x_1 \exists y_1^{\text{int}} \dots \forall x_n \exists y_n^{\text{int}} (f(x_1, y_1, \dots, x_n, y_n) \neq 0)$  is realized

by a proof-like term that depends only on  $n$ .

This theorem shows once again that any true arithmetical formula is realized.

---

For  $n = 1$ , the proof is very simple :

**Theorem.** Let  $\theta \in \text{QP}$  be such that  $\theta \star \underline{n} \cdot \xi \cdot \pi > \xi \star \underline{n} \cdot \theta \underline{n}^+ \xi \cdot \pi$  with  $\underline{n}^+ = (s)\underline{n}$ .

Then  $\theta \underline{0} \Vdash \forall x \left( \forall y^{\text{int}} (f(x, y) \neq 0 \rightarrow \perp) \rightarrow \perp \right)$

for every  $f : \mathbb{N}^2 \rightarrow 2$  such that  $\mathbb{N} \models \forall x \exists y (f(x, y) = 1)$ .

We simply need to prove  $\theta \underline{0} \Vdash \forall y^{\text{int}} (f(y) \neq 0 \rightarrow \perp) \rightarrow \perp$

for every  $f : \mathbb{N} \rightarrow 2$  such that  $\mathbb{N} \models \exists y (f(y) = 1)$ .

**Lemma.** Let  $\xi \Vdash \forall y^{\text{int}} (f(y) \neq 0 \rightarrow \perp)$  ; if  $\theta \underline{n} \xi \not\Vdash \perp$ , then  $f(n) = 0$  and  $\theta \underline{n}^+ \xi \not\Vdash \perp$ .

We have  $\theta \star \underline{n} \cdot \xi \cdot \pi \notin \perp$ , thus  $\xi \star \underline{n} \cdot \theta \underline{n}^+ \xi \cdot \pi \notin \perp$  ;

therefore  $\theta \underline{n}^+ \xi \not\Vdash f(n) \neq 0$  hence the result. QED

Suppose  $\theta \star \underline{0} \cdot \xi \cdot \pi \notin \perp$  ; the lemma gives  $f(n) = 0$  for all  $n \in \mathbb{N}$ , a contradiction. QED

---

We consider now the case  $n = 2$ , which is typical for the general case.

**Theorem.** Let  $\theta_x = \lambda t \lambda \sigma \lambda m \lambda n (x m) \lambda y (H \sigma m y n) ((t) (\Sigma) \sigma m y) m' n'$  ( $x$  is free) where  $H, \Sigma$  are closed  $\lambda$ -terms defined below ;  $\langle m', n' \rangle$  is the successor of  $\langle m, n \rangle$  in  $\mathbb{N}^2$ .

Then, for every  $f : \mathbb{N}^3 \rightarrow \{0, 1\}$ , there exists  $\phi : \mathbb{N} \rightarrow \mathbb{N}$  such that :

$\lambda x (Y) \theta_x 000 \Vdash \forall x^{\text{int}} \exists y \forall z^{\text{int}} (f(x, y, z) = 1) \rightarrow \forall x \forall z (f(x, \phi x, z) \neq 0)$ .

**Definition of  $H, \Sigma$ .** The variables  $m, n$  represent integers ;  $\eta$  an arbitrary term ; the variable  $\sigma$  represents a finite sequence of ordered pairs  $\langle m, \eta \rangle$ .

If no pair  $\langle m, \cdot \rangle$  is in  $\sigma$ , put  $\Sigma \sigma m \eta = \sigma \cup \langle m, \eta \rangle$ ,  $H \sigma m \eta = \eta$ .

Else, put  $\Sigma \sigma m \eta = \sigma$  ;  $H \sigma m \eta = \zeta$  for the first  $\langle m, \zeta \rangle$  appearing in  $\sigma$ .

**Proof by contradiction.** Suppose  $\xi \Vdash \forall x^{\text{int}} \left( \forall y \left[ \forall z^{\text{int}} (f(x, y, z) = 1) \rightarrow \perp \right] \rightarrow \perp \right)$  ;

$Y \theta_\xi 000 \not\Vdash \perp$  and  $f(x_0, \phi x_0, z_0) = 0$ .

We show, by recurrence on  $\langle m, n \rangle \leq \langle x_0, z_0 \rangle$ , that  $Y \theta_\xi \sigma_{mn} m n \not\Vdash \perp$ ,

with  $\sigma_{mn}, \eta_{mn}, b_{mn}$  defined by recurrence ; it's true for  $\sigma_{00} = 0$ . If it's true for  $\langle m, n \rangle$

we have  $Y \theta_\xi \sigma_{mn} m n \star \pi \notin \perp$ , i.e.  $\theta_\xi \star Y \theta_\xi \cdot \sigma_{mn} \cdot m \cdot n \cdot \pi \notin \perp$ , or else :

---

$\xi \star m \cdot \lambda y(H\sigma_{mn}myn)((Y\theta_\xi)(\Sigma)\sigma_{mn}my)m'n' \cdot \pi \notin \perp$ . Thus, there exists  $b_{mn}$  s.t. :  
 $\lambda y(H\sigma_{mn}myn)((Y\theta_\xi)(\Sigma)\sigma_{mn}my)m'n' \not\vdash \forall z^{\text{int}}(f(m, b_{mn}, z) = 1) \rightarrow \perp$  and thus  
there exists  $\eta_{mn} \Vdash \forall z^{\text{int}}(f(m, b_{mn}, z) = 1)$  such that

(\*)  $H\sigma_{mn}m\eta_{mn} \star n \cdot ((Y\theta_\xi)(\Sigma)\sigma_{mn}m\eta_{mn})m'n' \cdot \pi \notin \perp$ .

**Definition of  $\phi m$**  : i) if no pair  $\langle m, \cdot \rangle$  appears in  $\sigma_{mn}$  then put  $\phi m = b_{mn}$  ;

ii) else,  $\langle m, \phi m \rangle$  is the first (indeed only) pair  $\langle m, \cdot \rangle$  appearing in  $\sigma_{mn}$ .

Now, we have  $H\sigma_{mn}m\eta_{mn} \Vdash \forall z^{\text{int}}(f(m, \phi m, z) = 1)$  because :

in case (i)  $H\sigma_{mn}m\eta_{mn} = \eta_{mn}$  and  $\phi m = b_{mn}$  ; in case (ii), by induction on  $\langle m, n \rangle$

since  $H\sigma_{mn}m\eta_{mn} = \eta_{mq}$  with  $\langle m, q \rangle$  strictly before  $\langle m, n \rangle$ .

Thus  $H\sigma_{mn}m\eta_{mn}n \Vdash f(m, \phi m, n) \neq 1 \rightarrow \perp$ .

Now, we put  $\sigma_{m'n'} = (\Sigma)\sigma_{mn}m\eta_{mn}$ .

Thus, by (\*), we have  $Y\theta_\xi\sigma_{m'n'}m'n' \not\vdash f(m, \phi m, n) \neq 1$ .

therefore  $f(m, \phi m, n) = 1$  and  $Y\theta_\xi\sigma_{m'n'}m'n' \Vdash \perp$ .

Since  $f(x_0, \phi x_0, z_0) = 0$ , we have a contradiction if  $\langle m, n \rangle = \langle x_0, z_0 \rangle$ .

Else, we have done the recurrence step.

QED

---

Consider now a function  $f : \mathbb{N}^4 \rightarrow \{0, 1\}$  s.t.  $\mathbb{N} \models \forall u \exists x \forall y \exists z (f(u, x, y, z) = 0)$ .

This gives  $\forall u (\forall x \exists y \forall z (f(u, x, y, z) \neq 0) \rightarrow \perp)$ .

Thus, for every  $u \in \mathbb{N}$  and  $\phi : \mathbb{N} \rightarrow \mathbb{N}$  we get :

$$\|\forall x \forall z (f(u, x, \phi x, z) \neq 0)\| = \|\perp\| = \Pi.$$

It follows from the previous theorem that

$$\lambda x(Y)\theta_x 000 \Vdash \forall u (\forall x^{\text{int}} \exists y \forall z^{\text{int}} (f(u, x, y, z) = 1) \rightarrow \perp)$$

which is the case  $n = 2$  for arithmetical formulas.

---

## ⌋2 trivial

Let  $\delta$  be a proof-like term s.t.  $\delta \Vdash \forall x \neg \lceil 2 (x \neq 0, x \neq 1 \rightarrow \perp)$  (i.e.  $\lceil 2$  is trivial).

We have  $\delta \in |\top, \perp \rightarrow \perp| \cap |\perp, \top \rightarrow \perp|$ . Let  $\delta' = \lambda x \lambda y c c \lambda k (\delta) ((k)x)(k)y$ ; then

$$\xi \star \pi \in \perp \text{ or } \eta \star \pi \in \perp \Rightarrow \delta' \star \xi \cdot \eta \cdot \pi \in \perp$$

Thus,  $\delta' \Vdash X, Y \rightarrow X$  and  $\delta' \Vdash X, Y \rightarrow Y$  for every truth values  $X, Y$ .

**Theorem.**  $(\exists \Phi \in \text{QP})(\forall \theta \in \text{QP})(\forall X \subset \Pi)(\theta \Vdash X \Rightarrow \Phi \Vdash X)$ .

Define  $e$  (read *eval*) by the following program :

$e_{\underline{0}} = B, e_{\underline{1}} = C, e_{\underline{2}} = E, e_{\underline{3}} = I, e_{\underline{4}} = K, e_{\underline{5}} = W, e_{\underline{6}} = cc, e_{\underline{7}} = \delta$  ;

$e_{\underline{n+8}} = ((e)(p_0)\underline{n})(e)(p_1)\underline{n}$  ;

where  $p_0, p_1$  define a recursive bijection from  $\mathbb{N}$  onto  $\mathbb{N}^2$ .

For every  $\theta \in \text{QP}$ , there is an integer  $n$  s.t.  $e_{\underline{n}} > \theta$ .

Now define  $\phi$  by :  $\phi \star n \cdot \pi > \delta' \star e_{\underline{n}} \cdot (\phi)(s)\underline{n} \cdot \pi$ . Finally  $\Phi$  is  $\phi_{\underline{0}}$ .

Let  $\theta \in \text{QP}$  s.t.  $\theta \Vdash X$  ; thus, we have  $\phi_{\underline{n}} \Vdash X$  for some  $n$ ,

then  $\phi_{\underline{n-1}} \Vdash X, \dots$  ; eventually  $\phi_{\underline{0}} \Vdash X$ .

QED



---

## ⌋2 trivial

Let  $\mathcal{B} = \mathcal{P}(\Pi)$  be the Boolean algebra of truth values.

The order is defined by  $A \leq B \Leftrightarrow (\exists \theta \in \text{QP})(\theta \Vdash A \rightarrow B)$ .

Thus, the order on  $\mathcal{B}$  is defined by  $A \leq B \Leftrightarrow \Phi \Vdash A \rightarrow B$ .

**Theorem.**  $\mathcal{B}$  is a complete Boolean algebra :

If  $B_i (i \in I)$  is a family of truth values, then  $\inf_{i \in I} B_i = \bigcup_{i \in I} B_i$ .

Let  $A \leq B_i$  for  $i \in I$ . Then  $\Phi \Vdash A \rightarrow B_i$ , thus  $\Phi \Vdash A \rightarrow \bigcup_{i \in I} B_i$ .

Conversely  $I \Vdash \bigcup_{i \in I} B_i \rightarrow B_{i_0}$ .

QED

Thus, the realizability model is, in fact, a *forcing model*.

The converse is also true : in the case of forcing, the realizability algebra is a commutative idempotent monoid with a unity  $\mathbf{1}$  ; then  $\text{QP} = \{\mathbf{1}\}$ .

We have  $\mathbf{1} \Vdash X, Y \rightarrow X$  and  $X, Y \rightarrow Y$  ; thus ⌋2 is trivial.

---

## $\mathbb{J}2$ with 4 elements

Let  $d$  be a term constant and suppose  $\perp$  has the following property :

*If two out of the three processes  $\xi \star \pi, \eta \star \pi, \zeta \star \pi$  are in  $\perp$  then  $d \star \xi \cdot \eta \cdot \zeta \cdot \pi \in \perp$ .*

We have  $d \in |\top, \perp, \perp \rightarrow \perp| \cap |\perp, \top, \perp \rightarrow \perp| \cap |\perp, \perp, \top \rightarrow \perp|$ .

Thus  $d \Vdash \forall x \mathbb{J}2 \forall y \mathbb{J}2 (x \neq 0, y \neq 1, x \neq y \rightarrow xy \neq x)$

i.e.  $d \Vdash \text{"}\mathbb{J}2 \text{ has at most 4 elements"}$

We now build a model in which  $\mathbb{J}2$  has exactly 4 elements.

The only term constants are the elementary combinators,  $cc$  and  $d$ .

There are two stack constants  $\pi^0, \pi^1$ . Let  $\omega = (WI)(W)I = (\lambda x xx)\lambda x xx$ .

For  $i \in \{0, 1\}$ , let  $\Lambda^i$  (resp.  $\Pi^i$ ) be the set of terms (resp. stacks)

which contain the only stack constant  $\pi^i$ .

For  $i, j \in \{0, 1\}$ , define  $\perp_j^i$  as the least set  $P \subset \Lambda^i \star \Pi^i$  of processes such that :

1.  $\omega \star \underline{j} \cdot \pi^i \in P$  ;
2.  $\xi \star \pi \in \Lambda^i \star \Pi^i, \xi \star \pi \succ \xi' \star \pi' \in P \Rightarrow \xi \star \pi \in P$  ( $P$  is saturated in  $\Lambda^i \star \Pi^i$ ) ;
3. if 2 out of the 3 processes  $\xi \star \pi, \eta \star \pi, \zeta \star \pi$  are in  $P$ , then  $d \star \xi \cdot \eta \cdot \zeta \cdot \pi \in P$ .

---

We define  $\perp$  by :  $\Lambda \star \Pi \setminus \perp = \bigcup_{i \in \{0,1\}} (\Lambda^i \star \Pi^i \setminus \perp_j^i)$

In other words, a process is in  $\perp$  iff

either it is in  $\perp_0^0 \cup \perp_1^1$  or it contains both stack constants  $\pi^0, \pi^1$ .

**Lemma.** If  $\xi \star \pi \in \perp_j^i$  and  $\xi \star \pi > \xi' \star \pi'$  then  $\xi' \star \pi' \in \perp_j^i$  (closure by reduction).

Suppose  $\xi_0 \star \pi_0 > \xi'_0 \star \pi'_0$ ;  $\xi_0 \star \pi_0 \in \perp_j^i$ ;  $\xi'_0 \star \pi'_0 \notin \perp_j^i$ ;

We may suppose that  $\xi_0 \star \pi_0 > \xi'_0 \star \pi'_0$  is exactly one step.

Then  $\perp_j^i \setminus \{\xi_0 \star \pi_0\}$  has properties 1,2,3 defining  $\perp_j^i$ ; contradiction. QED

**Lemma.**  $\perp_0^i \cap \perp_1^i = \emptyset$ .

We prove that  $\Lambda^i \star \Pi^i \setminus \perp_1^i \supset \perp_0^i$  by showing properties 1, 2, 3.

1.  $\omega \star \underline{0} \cdot \pi^i \notin \perp_1^i$  because  $\perp_1^i \setminus \{\omega \star \underline{0} \cdot \pi^i\}$  has properties 1, 2, 3 defining  $\perp_1^i$ .

2. Follows from previous lemma.

3. Suppose  $\xi \star \pi, \eta \star \pi \notin \perp_1^i$ ; then  $d \star \xi \cdot \eta \cdot \zeta \cdot \pi \notin \perp_1^i$

because  $\perp_1^i \setminus \{d \star \xi \cdot \eta \cdot \zeta \cdot \pi\}$  has properties 1, 2, 3 defining  $\perp_1^i$ . QED

**Theorem.** This realizability model is coherent.

Let  $\theta \in \text{QP}$  s.t.  $\theta \star \pi^0 \in \perp_0^0$  and  $\theta \star \pi^1 \in \perp_1^1$ . Then  $\theta \star \pi^0 \in \perp_0^0 \cap \perp_1^0$ . QED

---

**Remark.** If  $\pi \in \Pi \setminus (\Pi_0 \cup \Pi_1)$ , then  $\xi \star \pi \in \perp$  for every term  $\xi$ .

Thus, we can remove these stacks and consider only  $\Pi^0 \cup \Pi^1$ .

We define two individuals in this realizability model :

$$\gamma_0 = (\{0\} \times \Pi^0) \cup (\{1\} \times \Pi^1) ; \gamma_1 = (\{1\} \times \Pi^0) \cup (\{0\} \times \Pi^1).$$

Obviously,  $\gamma_0, \gamma_1 \subset \mathbb{J}2 = \{0, 1\} \times \Pi$ . Now we have :

$$\|\forall x(x \notin \gamma_0)\| = \Pi^0 \cup \Pi^1 = \|\perp\| ; \omega_{\underline{0}} \Vdash 0 \notin \gamma_0 \text{ et } \omega_{\underline{1}} \Vdash 1 \notin \gamma_0.$$

It follows that  $\gamma_0$  is not  $\varepsilon$ -empty and that every  $\varepsilon$ -element of  $\gamma_0$  is  $\neq 0, 1$ .

Thus the Boolean algebra  $\mathbb{J}2$  is not trivial and has exactly 4  $\varepsilon$ -elements.

We have  $\xi \Vdash \forall x \mathbb{J}2(x \varepsilon \gamma_0, x \varepsilon \gamma_1 \rightarrow \perp)$  for every term  $\xi$  :

Indeed,  $|i \varepsilon \gamma_0| = \{k_\pi ; \pi \in \Pi^i\}$  for  $i = 0, 1$  and  $\xi \star k_{\rho_0} \cdot k_{\rho_1} \cdot \pi \in \perp$  if  $\rho_i \in \Pi^i$ .

It follows that  $\gamma_0, \gamma_1$  are the singletons of the  $\varepsilon$ -elements  $\neq 0, 1$  of  $\mathbb{J}2$ .

**Remark.** We can easily modify this construction in order to obtain for  $\mathbb{J}2$  any finite Boolean algebra.

---

## Denotational semantics

T. Ehrhard has found a method which converts usual models of  $\lambda$ -calculus into realizability algebras, by defining stacks, cc and  $k_\pi$  in such models.

The construction of stacks was also given by T. Streicher.

We need to avoid *parallel or*, because we don't want to get *forcing models*.

Thus, our example will be the simplest *coherent model of  $\lambda$ -calculus*.

Let us recall (one of) its construction.

Let  $\mathbf{o}$  be a fixed set which is not an ordered pair.

The the set  $V$  of *formulas* is the smallest set s.t. :

$\mathbf{o} \in V$  ; if  $\alpha \in V$ ,  $a \in \mathcal{P}_f(V)$  and  $\langle a, \alpha \rangle \neq \langle \emptyset, \mathbf{o} \rangle$  then  $\langle a, \alpha \rangle \in V$

( $\mathcal{P}_f(V)$  is the set of finite subsets of  $D$ ).

If  $a \in \mathcal{P}_f(V)$  and  $\alpha \in V$ , we set  $a \rightarrow \alpha = \langle a, \alpha \rangle$  except that  $(\emptyset \rightarrow \mathbf{o}) = \mathbf{o}$ .

Every element of  $V$  except  $\mathbf{o}$  is an ordered pair.

If  $\alpha \in V$ , its rank  $r(\alpha)$  is the total number of  $\rightarrow$  in  $\alpha$ .

---

Each  $\alpha \in V$  has a unique normal form  $\alpha = (a_1, \dots, a_k \rightarrow \circ)$

with  $k \in \mathbb{N}$ ,  $a_1, \dots, a_k \in \mathcal{P}_f(V)$  and  $a_k \neq \emptyset$ . Then  $\alpha = (a_1, \dots, a_k, \emptyset, \dots, \emptyset \rightarrow \circ)$ .

The *truth value*  $|\alpha| \in \{0, 1\}$  of a formula  $\alpha$  is defined by induction :

$|a_1, \dots, a_k \rightarrow \circ| = 1$  iff  $\exists i (\exists \beta \in a_i) (|\beta| = 0)$ .

If  $\alpha = (a_1, \dots, a_k \rightarrow \circ)$ ,  $\beta = (b_1, \dots, b_k \rightarrow \circ)$  we define

$$\alpha \sqcap \beta = (a_1 \cup b_1, \dots, a_k \cup b_k \rightarrow \circ).$$

This operation is associative, commutative and idempotent ;  $\circ$  is neutral ;

it defines an order relation :  $\alpha \leq \beta \Leftrightarrow b_1 \subset a_1, \dots, b_k \subset a_k$ .

Define a subset  $D$  of  $V$  (the *web*) by induction on the rank :

$(a_1, \dots, a_k \rightarrow \circ) \in D$  iff, for  $1 \leq i \leq k$ ,

$a_i \subset D$  and  $(\forall \beta, \gamma \in a_i) (\beta \neq \gamma \Rightarrow \beta \sqcap \gamma \notin D)$  ( $a_i$  is an *antichain* of  $D$ ).

*D is a final segment of V* : let  $\alpha = (a_1, \dots, a_k \rightarrow \circ)$ ,  $\beta = (b_1, \dots, b_k \rightarrow \circ)$ ,

$\alpha \in D, \alpha \leq \beta$ . Then  $b_i \subset a_i$  and  $a_i$  is an antichain of  $D$ , thus so is  $b_i$ .

$\alpha, \beta \in D$  are called *compatible* if  $\alpha \sqcap \beta \in D$  ; in symbols  $\alpha \asymp \beta$ .

If  $\alpha_1, \dots, \alpha_n$  are pairwise compatible, then  $\alpha_1 \sqcap \dots \sqcap \alpha_n \in D$ .

---

## The realizability algebra

$\Lambda_D$  is the set  $\mathcal{A}(D)$  of antichains of  $D$ , i.e.  $t \subset D$  is a term iff  $(\forall \alpha, \beta \in t)(\alpha \sqcap \beta \in D \rightarrow \alpha = \beta)$ .

$\Pi_D$  is the set  $\mathcal{S}(D)$  of filters of  $D$ , i.e.  $\pi \subset D$  is a stack iff  $(\forall \alpha, \beta \in \pi) \alpha \sqcap \beta \in \pi ; \forall \alpha \forall \beta (\alpha \in \pi, \alpha \leq \beta \rightarrow \beta \in \pi) ; \mathbf{0} \in \pi$ .

**Remark.**  $\Pi_D$  can be identified with  $\Lambda_D^{\mathbb{N}}$ : a sequence of terms  $t_n (n \in \mathbb{N})$  correspond with the filter  $\{a_1, \dots, a_k \rightarrow \mathbf{0} ; a_1 \subset t_1, \dots, a_k \subset t_k\}$ .

$\Lambda_D \star \Pi_D$  is  $\{0, 1\}$  and  $\perp$  is  $\{1\}$ .

If  $t \in \Lambda_D, \pi \in \Pi_D$  then  $t \star \pi \in \perp$  iff  $t \cap \pi \neq \emptyset$  (i.e.  $t \cap \pi$  is a singleton).

$t \bullet \pi = \{a \rightarrow \alpha ; a \subset t, \alpha \in \pi\} ;$

$tu = \{\alpha \in D ; (\exists a \subset u)(a \rightarrow \alpha) \in t\} ;$

$K$  is the set of all formulas :  $\{\alpha\}, \emptyset \rightarrow \alpha$  for  $\alpha \in D$ .

$S$  is the set of all formulas :

$\{a_0, \{\alpha_1, \dots, \alpha_k\} \rightarrow \beta\}, \{a_1 \rightarrow \alpha_1, \dots, a_k \rightarrow \alpha_k\}, a_0 \cup a_1 \cup \dots \cup a_k \rightarrow \beta$

with  $\{\alpha_1, \dots, \alpha_k\} \in \mathcal{A}(D)$  and  $a_0 \cup a_1 \cup \dots \cup a_k \in \mathcal{A}(D)$ .

---

$k_\pi$  is the set of formulas :  $(\{\alpha\} \rightarrow o)$  for  $\alpha \in \pi$  ;

$cc$  is the set of all formulas :

$\{a \rightarrow \alpha\} \rightarrow \alpha \sqcap \alpha_1 \sqcap \dots \sqcap \alpha_k$  with  $a = \{\{\alpha_1\} \rightarrow o, \dots, \{\alpha_k\} \rightarrow o\}$  and  $\alpha \sqcap \alpha_1 \sqcap \dots \sqcap \alpha_k \in D$ .

QP is defined as the set of  $t \in \Lambda_D$  s.t.  $|t| = 1$  i.e.  $(\forall \alpha \in t)(|\alpha| = 1)$ .

We have  $K, S, cc \in QP$  ;  $t, u \in QP \Rightarrow tu \in QP$ .

The model is *coherent* because  $|t| = 1 \Rightarrow o \notin t$  i.e.  $t \star \{o\} \notin \perp$ .

**Lemma 1.**  $t \Vdash \top, \dots, \top \rightarrow \perp$  iff  $t = \{o\}$ .

Indeed,  $t \star \emptyset \dots \emptyset \cdot \{o\} \in \perp \Rightarrow t = \{o\}$

QED

**Lemma 2.** If  $t \in |\top, \perp \rightarrow \perp| \cap |\perp, \top \rightarrow \perp|$  then  $t = \{o\}$ .

We have  $t \cap \emptyset \cdot \{o\} \cdot \{o\} \neq \emptyset$  and  $t \cap \{o\} \cdot \emptyset \cdot \{o\} \neq \emptyset$  ; thus

$(\emptyset, a \rightarrow o) \in t$  and  $(b, \emptyset \rightarrow o) \in t$  with  $a, b \subset \{o\}$ .

These two formulas are compatible and therefore equal ; thus  $a = b = \emptyset$ .

QED

It follows that  $\perp \Vdash |\top, \perp \rightarrow \perp| \cap |\perp, \top \rightarrow \perp| \rightarrow \perp$  i.e.

$\perp \Vdash \forall x \mathbb{I}^2(x \neq 0, x \neq 1 \rightarrow \perp) \rightarrow \perp$ . Therefore :

*The Boolean algebra  $\mathbb{I}^2$  is non trivial.*



---

**Lemma 3.** If  $u \Vdash \perp, \perp \rightarrow \perp$  then  $u$  contains one of the formulas :

$o ; \{o\} \rightarrow o ; \emptyset, \{o\} \rightarrow o ; \{o\}, \{o\} \rightarrow o.$

We have  $u \cap \{o\} \cdot \{o\} \cdot \{o\} \neq \emptyset$ , thus there exist  $a, b \subset \{o\}$  s.t.  $(a, b \rightarrow o) \in u.$

QED

**Lemma 4.** Let  $t \in \Lambda_D$  contain the 4 incompatible formulas :

$\{o\} \rightarrow o ; \{\{o\} \rightarrow o\}, \{o\} \rightarrow o ; \{\emptyset, \{o\} \rightarrow o\}, \{o\} \rightarrow o ; \{\{o\}, \{o\} \rightarrow o\}, \{o\} \rightarrow o.$

Then  $t \Vdash |\top, \perp \rightarrow \perp| \cap |\perp, \top \rightarrow \perp|, \top \rightarrow \perp$  and  $t \Vdash (\perp, \perp \rightarrow \perp), \perp \rightarrow \perp.$

By lemma 2, the first conclusion is  $t \Vdash \perp \rightarrow \perp$  ; it is satisfied because  $(\{o\} \rightarrow o) \in t.$

Now, let  $u \Vdash \perp, \perp \rightarrow \perp$  ; we must show  $t \cap u \cdot \{o\} \cdot \{o\} \neq \emptyset$

which follows immediately from lemma 3.

QED

**Theorem.** The Boolean algebra  $\mathbb{J}2$  is atomless.

We have  $t \Vdash \forall x \mathbb{J}2 \left( \forall y \mathbb{J}2 (xy \neq 0, xy \neq x \rightarrow \perp), x \neq 0 \rightarrow \perp \right)$  iff

$t \Vdash |\top, \perp \rightarrow \perp| \cap |\perp, \top \rightarrow \perp|, \top \rightarrow \perp$  and  $t \Vdash (\perp, \perp \rightarrow \perp), \perp \rightarrow \perp.$

Hence the result by lemma 4.

QED

---

## Integers

In the sequel, we use truth values defined by subsets  $|V|$  of  $\Lambda$ . They may be used in formulas only before a  $\rightarrow$ .

If  $|V| \subset \Lambda$ ,  $\|A\| \subset \Pi$ , we define  $\|V \rightarrow A\| = \{t \cdot \pi; t \in |V|, \pi \in \|A\|\}$ .

In particular  $\|\neg V\| = \{t \cdot \pi; t \in |V|, \pi \in \Pi\}$ .

**Lemma 5.** If  $(\forall t \in \Lambda)(t \in |V| \Rightarrow \theta t \in |W|)$  then  $\lambda x x \circ \theta \Vdash \neg W \rightarrow \neg V$ .

We shall sometimes write  $\theta \Vdash V \rightarrow W$  in such a case.

Now, define the formulas :

$v_0 = (\{0\} \rightarrow 0)$  ;  $v_1 = (\emptyset, \{0\} \rightarrow 0)$  ; ... ;  $v_n = (\emptyset, \dots, \emptyset, \{0\} \rightarrow 0)$  ; ... ;

and the terms  $\bar{n} = \{v_n\}$  ;  $\text{suc} = \{(\{v_0\} \rightarrow v_1), \dots, (\{v_i\} \rightarrow v_{i+1}), \dots\}$ .

Define the unary predicate  $N$  by :

$|Nn| = \{\bar{n}\}$  if  $n \in \mathbb{N}$  ;  $|Nn| = \emptyset$  if  $n \notin \mathbb{N}$ .

Then we have easily  $\lambda x(x) \bar{0} \Vdash \neg \neg N0$  ;  $\text{suc} \Vdash Nn \rightarrow N(n+1)$  for every  $n$  ;

i.e.  $\lambda x x \circ \text{suc} \Vdash \forall x(\neg N(x+1) \rightarrow \neg Nx)$ .

We have shown :  $\Vdash \forall x^{\text{int}} \neg \neg Nx$ .

---

**Theorem 6.** Let  $u_n (n \in \mathbb{N})$  be any sequence of terms and define :

$\theta = \{(\{v_n\} \rightarrow \alpha) ; n \in \mathbb{N}, \alpha \in u_n\}$ . Then  $\theta \bar{n} = u_n$  for all  $n \in \mathbb{N}$ .

If every  $u_n$  is in QP, then  $\theta \in \text{QP}$ .

We show that  $\theta \in \Lambda_D$  : if  $(\{v_m\} \rightarrow \alpha) \simeq (\{v_n\} \rightarrow \beta)$  then  $\{v_m, v_n\}$  is an antichain and therefore  $m = n$  ; thus  $\alpha, \beta \in u_m$  ; but  $\alpha \simeq \beta$  and therefore  $\alpha = \beta$ .

$\theta\{v_n\} = u_n$  is obvious.

QED

Define the unary predicate  $\text{ent}(x)$  by :

$|\text{ent}(n)| = \{\underline{n}\}$  (Church integer) for  $n \in \mathbb{N}$  ;  $|\text{ent}(n)| = \emptyset$  if  $n \notin \mathbb{N}$ .

We already know (general theory) that  $\text{ent}(x)$  is equivalent to  $\text{int}(x)$ .

Apply lemma 5 and theorem 6 above with  $u_n = \{\underline{n}\}$ .

This gives  $\theta \Vdash Nn \rightarrow \text{ent}(n)$  and therefore  $\lambda x x \circ \theta \Vdash \forall x (\neg \text{ent}(x) \rightarrow \neg Nx)$ .

Finally we have shown that the predicates  $Nx, \text{int}(x), \text{ent}(x)$  are equivalent.

In the following, we use  $Nx$  which is the simplest.

---

**Corollary.** If  $\theta_n \Vdash F(n)$ , with  $\theta_n \in \text{QP}$  for all  $n \in \mathbb{N}$ , then there exists  $\theta \in \text{QP}$  s.t.  $\theta \Vdash \forall n^{\text{int}} F(n)$ .

Applying theorem 6, we get  $\theta_{\underline{n}} \Vdash F(n)$  for all  $n \in \mathbb{N}$ , thus  $\theta \Vdash \forall n^{\text{int}} F(n)$ . QED

By the above corollary, the set of formulas which are realized by a proof-like term is closed by the  $\omega$ -rule.

Thus there exists a realizability model *which is an  $\omega$ -model*.

Let  $\mathcal{B} = \mathcal{P}(\Pi)$  be the Boolean algebra of truth values.

The order is defined by  $\|A\| \leq \|B\| \Leftrightarrow (\exists \theta \in \text{QP})(\theta \Vdash A \rightarrow B)$ .

**Theorem.**  $\mathcal{B}$  is a countably complete Boolean algebra :

If  $\|B(n)\|_{n \in \mathbb{N}}$  is a sequence of truth values, then  $\inf_{n \in \mathbb{N}} \|B(n)\| = \|\forall x^{\text{int}} B(x)\|$ .

Let  $\|A\| \leq \|B(n)\|$  for every  $n \in \mathbb{N}$ . Then  $\theta_n \Vdash A \rightarrow B(n)$  for some sequence  $\theta_n \in \text{QP}$ .

By the previous corollary, we get  $\theta \Vdash \|A \rightarrow \forall x^{\text{int}} B(x)\|$  i.e.  $\|A\| \leq \|\forall x^{\text{int}} B(x)\|$ .

Conversely,  $\|\forall x^{\text{int}} B(x)\| \leq \|B(n)\|$  because  $\lambda x(x) \underline{n} \Vdash \forall x^{\text{int}} B(x) \rightarrow B(n)$ . QED